

A search for radio pulsars around low-mass white dwarfs

Joeri van Leeuwen*, Robert D. Ferdman, Sol Meyer and Ingrid Stairs

Department of Physics & Astronomy, University of British Columbia, 6224 Agricultural Road, Vancouver B.C. V6T 1Z1, Canada.

Accepted Received

ABSTRACT

Low-mass white dwarfs can either be produced in low-mass X-ray binaries by stable mass transfer to a neutron star, or in a common-envelope phase with a heavier white dwarf companion. We have searched 8 low-mass white dwarf candidates recently identified in the Sloan Digital Sky Survey for radio pulsations from pulsar companions, using the Green Bank Telescope at 340MHz. We have found no pulsations down to flux densities of $0.6\text{--}0.8\text{ mJy kpc}^{-2}$ and conclude that a given low-mass helium-core white dwarf has a probability of $< 0.18 \pm 0.05$ of being in a binary with a radio pulsar.

Key words: pulsars: general — white dwarfs

1 INTRODUCTION

High-precision timing of millisecond and binary pulsars has applications ranging from setting limits on the stochastic gravitational-wave background (e.g. Jaffe & Backer 2003), to studying binary evolution, neutron-star masses and equations of state (e.g. Lattimer & Prakash 2001), and testing the predictions of general relativity (e.g. Taylor & Weisberg 1989; Lyne et al. 2004). Wide-area systematic searches have historically been very successful in finding millisecond pulsars (e.g. Lorimer et al. 1996). In contrast to these wide-area surveys, directed pulsar searches use typical associations of neutron stars with other objects to increase the chance of finding a pulsar in a given telescope pointing. This allows direct searches to use fewer pointings, with longer integration times and hence better sensitivity. Directed searches towards certain object types also elucidate their evolutionary relation with radio pulsars through the incidence of the associations (cf. Staelin & Reifenstein 1968). The classic example is searching for young pulsars toward pulsar wind nebulae and supernova remnants, often with known central X-ray point sources (e.g. Camilo et al. 2002), and the success of these searches has enabled meaningful statistics on the interactions between the radio pulsars and the supernova remnants (e.g. Chevalier 2005). Directed searches for recycled *millisecond* pulsars have so-far been limited to globular clusters (e.g. Ransom et al. 2005) and a few isolated steep-spectrum targets (e.g. Backer et al. 1982; Navarro et al. 1995).

The situation has changed with the publication of catalogs of white dwarfs from deep surveys. In particular, Kleinman et al. (2004) list over 2500 spectroscopically




Figure 1. Measured temperatures and surface gravities plotted onto simulated evolutionary tracks for low-mass helium-core white dwarfs of different mass. Shown are the 7 candidates (open circles) and 2 pulsar companions (filled squares) that fall within the simulated range. Data taken from Kleinman et al. (2004), Eisenstein et al. (2006) and Bassa et al. (2006), plotted over evolutionary tracks from Althaus et al. (2001) and Serenelli et al. (2001) as cited in Liebert et al. (2004).

identified white dwarfs from the first data release of the Sloan Digital Sky Survey (SDSS, York et al. 2000). Among these are a handful which appear to have very low surface gravities ($\log g \leq 7.5$) and fairly high effective temperatures of $2\text{--}3 \times 10^4\text{ K}$, indicating probable masses of less than $0.45 M_{\odot}$ and a helium composition (Figure 1). These objects are described in more detail in Liebert et al. (2004), in which one object, SDSS J123410.37–022802.9, is shown to have an excellently determined spectrum and a likely

* e-mail: joeri@astro.ubc.ca

Name	Integration time (s)	Flux limit (mJy)
SDSS J002207.65−101423.5	1400 + 2800	0.16
SDSS J081136.34+461156.4	1400 + 2800	0.16
SDSS J102228.02+020035.2	1400 + 2800	0.13
SDSS J105611.03+653631.5	2 × 1400	0.21
SDSS J123410.37−022802.9	3 × 2800	0.19
SDSS J130422.65+012214.2	3 × 1400	0.24
SDSS J142601.48+010000.2	1400 + 2800	0.17
SDSS J234536.48−010204.8	2 × 2800	0.15

Table 1. Upper limits were derived using the longest individual integration time, a 22 K receiver temperature, sky temperatures from Haslam et al. (1981) using a spectral index of -2.6 , a minimum detection SNR of 10, a 2.0 K Jy^{-1} telescope gain and a constant 10% pulsar duty cycle.

mass of $\sim 0.18\text{--}0.19 M_{\odot}$. Such a low-mass, helium-core white dwarf is the prototypical companion of a highly recycled millisecond pulsar: in the low-mass X-ray binary phase much of the white-dwarf envelope is transferred to the neutron star, spinning it up. While this is not the only plausible explanation for the existence of this star (a white-dwarf companion is also a potential explanation), the match with the masses of known pulsar companions such as those of PSRs J1012+5307 ($0.16 \pm 0.02 M_{\odot}$, Callanan et al. 1998), J0437−4715 ($0.236 \pm 0.017 M_{\odot}$, van Straten et al. 2001), J0218+4232 ($0.20 \pm 0.02 M_{\odot}$, Bassa et al. 2003) and J1911−5958A ($0.175 \pm 0.010 M_{\odot}$, Bassa et al. 2006) is tantalizing.

Using the Green Bank Telescope (GBT) we have searched for radio pulsations from possible neutron-star companions to the 8 white dwarfs listed in Liebert et al. (2004) and Table 1. We describe our observing setup in § 2, our results in § 3 and discuss our findings in § 4.

2 OBSERVING SETUP

All sources were observed with the GBT on two or more occasions between October 21 and 26, 2004. At a central observing frequency of 340 MHz the Spigot backend (Kaplan et al. 2005) provided 50 MHz of bandwidth divided into 1024 spectral channels, recording total intensity in 16 bits every $81.92 \mu\text{s}$. Sessions started with 1-minute observations of bright test pulsar PSR J1012+5307 or PSR J2317+1439, followed by several 23-minute (2^{24} samples) or 46-minute (2^{25} samples) searches in the direction of the white dwarfs listed in Table 1.

Over the following months the data were processed using the SIGPROC package (Lorimer 2001) on the GASP computer cluster (Demorest et al. 2004). As described in more detail below, data were examined for radio interference, corrected for interstellar dispersion and binary acceleration, and searched for periodic signals.

Each observation was checked for interference by summing all frequency channels at zero dispersion measure (DM), searching for the strongest periodicities in the resulting time series and then localizing the matching interference within the data. Usually several affected frequency channels or data segments had to be removed, mostly containing non-




Figure 2. Detection signal-to-noise versus trial dispersion measure and acceleration for a partly simulated 40-minute observation of PSR J1012+5307. Using the PSR J1012+5307 binary parameters from Lange et al. (2001), we extrapolated one of our 1-minute test observations to a 40-minute observation around orbital phase 0.25. These data were subsequently processed in our search pipeline which detected the pulsar at an acceleration of 4.8 m s^{-2} and a DM of 9.0 pc cm^{-3} , close to the literature value of 9.0233 (Nicastro et al. 1995).

periodic bursts or (harmonics of) 1.266 and 60 Hz interference. The source of the former frequency is unclear.

We next formed multiple trial time series for DMs between 5 pc cm^{-3} and twice the maximum expected DM in the direction of each white dwarf, which is on average around 40 pc cm^{-3} (Cordes & Lazio 2002), using 0.1 to 0.5 pc cm^{-3} intervals.

We are specifically targeting millisecond pulsars in binary systems and the acceleration that accompanies the pulsar’s orbital motion can introduce significant changes in the apparent pulse period, hampering detection of a periodic pulse signal. The maximum acceleration of 90% of all currently known pulsars falls within the -25 m s^{-2} to $+25 \text{ m s}^{-2}$ range (Manchester et al. 2005) and we therefore resample each dedispersed time series for 250 trial accelerations throughout this range.

For each dedispersed and resampled time series we then compute the discrete Fourier transform and look for peaks, caused by periodic signals. As different pulse shapes and duty cycles produce peaks at different harmonics of the pulse frequency, we also fold the spectrum 2,4,8 and 16 times to add these harmonics. For each peak in these 5 raw and folded spectra that has a signal-to-noise ratio (SNR) exceeding 5σ , the signal frequency and SNR are stored.

After candidates have been identified in this fashion for all trial dedispersion and acceleration values, occurrences of the same period at different DMs and accelerations are




Figure 3. Estimated minimum detectable flux S_{min} as a function of period, for 4 characteristic dispersion measures (0, 20, 40 and 60 pc cm^{-3}). Curves are calculated for the observing setup described in § 2 and Table 1, for a 2800 s integration time and 30 K sky temperature.

combined to summarize and plot the signal properties. The first diagnostic is a two dimensional plot of SNR versus DM and acceleration, as shown in Figure 2 for test pulsar PSR J1012+5307. In order of peak SNR these plots are inspected for a well-defined response in both DM and acceleration. For the second diagnostic the time series are dedispersed, de-accelerated and folded at the best values for DM, acceleration and period. This allows us to check on pulse shapes, signal bandwidth and signal transience or persistence.

3 RESULTS

None of the observations were found to contain signals that merited follow-up. In Table 1 we estimate the pulsar-flux upper limits for the longest observation of each candidate. The upper limits quoted are for pulsars with periods of 20 ms or more, for which our search algorithm is most sensitive; in Figure 3 we show the typical dependence of this minimum detectable flux S_{min} on the pulsar period and dispersion measure.

4 DISCUSSION

There are several evolutionary scenarios thought to produce low-mass helium-core white dwarfs. In all these the progen-

itor of the observed white dwarf interacts with a close binary companion and loses its envelope. In one channel of evolution, the progenitor’s companion is a heavier white dwarf and a common-envelope phase exposes the progenitor’s helium core. As the companion white dwarf is older and cooler it is much dimmer and harder to detect than the observed white dwarf. In the other channel of evolution the progenitor of the observed white dwarf (WD) is in a binary with a neutron star (NS). As the progenitor evolves onto the red giant branch it overflows its Roche lobe and a phase of mass transfer starts that both recycles the pulsar and removes part of the progenitor’s hydrogen envelope (cf. Bhattacharya & van den Heuvel 1991).

We will outline the factors that determine radio pulsar detectability to estimate the probability P_{MSP} that a individual helium-core white dwarf was formed in the presence of a neutron star, as opposed to formation near another white dwarf.

We first evaluate our luminosity completeness. The combination of our radio-flux upper limits with the white-dwarf distance estimates allows us to constrain the radio-pulsar luminosities. Typically, our candidate white dwarfs are 3–4 magnitudes brighter in r than the companion to PSR J0218+4232 (18.2–20.5 versus 23.9, Eisenstein et al. 2006; Bassa et al. 2003) and hotter by a factor of 2. We therefore use the low end of the distance range to PSR J0218+4232 (2.5–4 kpc, Bassa et al. 2003) to estimate the distance for each candidate to be 2 kpc. At this distance our radio-flux upper limits amount to a luminosity of 0.6–0.8 mJy kpc^{-2} . As 95% of currently known field millisecond pulsars are brighter than this (Manchester et al. 2005) we assume we are complete to a level of $P_{lum}=0.95$.

Millisecond pulsars generally have relatively wide beams that show up as high duty cycles ($\sim 20\%$ on average) and increase the chance that the beam sweeps over the Earth. For millisecond pulsars this beaming fraction P_{beam} is thought to be as high as 0.7 ± 0.2 (Kramer et al. 1998).

The maximum acceleration of 90% of all currently known pulsars falls in our acceleration search range, and as the 10% with higher maximum accelerations are still detectable during large parts of their orbit, where the component of their motion in our line of sight is less, we estimate $P_{acc}=0.95$.

As more than half of millisecond pulsars with known characteristic ages appear to be older than 5 Gyr (Manchester et al. 2005) we assume recycled radio pulsars remain active over the cooling lifetime of white dwarfs.

We will use our non-detections and the luminosity, acceleration and beaming fractions mentioned above to estimate the probability P_{MSP} that a low-mass helium-core white dwarf is made through the channel involving a millisecond-pulsar companion. We assume that our $N=8$ candidates were correctly identified as low-mass helium white dwarfs. Taking radio frequency interference into account we estimate that our pulsar search-algorithm success rate P_{eff} is 80%.

In this case the assumption that our non-detection was the most probable outcome of our experiment:

$$(1 - P_{lum} \times P_{acc} \times P_{beam} \times P_{eff} \times P_{MSP})^N > \frac{1}{2} \quad (1)$$

leads us to conclude that $P_{MSP} < 0.18 \pm 0.05$.

Our upper limit for P_{MSP} is the first measurement of the probabilities in the formation of low-mass helium-core white dwarfs. In comparison, for general white dwarfs the mass-ratio distribution appears to peak around unity, strongly favoring WD-WD over WD-NS binaries: Pinsonneault & Stanek (2006) find that 50% of detached eclipsing binaries in the Small Magellanic Cloud have very similar masses and Maxted et al. (2002) show that WD-WD binary stars relatively often have mass ratios close to 1. Simulations by Schneider et al. (2001) predict the birth rate for general WD-WD binaries to be 100 times higher than for WD-NS binaries – a significant fraction of potential WD-NS binaries are disrupted in the supernova that creates the neutron star.

5 CONCLUSIONS

We have searched for radio pulsars around 8 low-mass white dwarf candidates from the SDSS and have found none down to a limit at the 5th percentile of the currently known millisecond-pulsar luminosity distribution. Low-mass helium-core white dwarfs can form around either a heavier white dwarf or a neutron star: we estimate the probability of formation through the latter evolution channel to be $< 0.18 \pm 0.05$.

6 ACKNOWLEDGEMENTS

Van Leeuwen and Ferdman wish to thank HIA for travel support. Stairs holds an NSERC UFA. Pulsar research at UBC is supported by a Discovery Grant. The National Radio Astronomy Observatory is a facility of the National Science Foundation operated under cooperative agreement by Associated Universities, Inc.

REFERENCES

- Althaus L. G., Serenelli A. M., Benvenuto O. G., 2001, *MNRAS*, 323, 471
- Backer D. C., Kulkarni S. R., Heiles C., Davis M. M., Goss W. M., 1982, *Nature*, 300, 615
- Bailes M., 1989, *ApJ*, 342, 917
- Bassa C. G., van Kerkwijk M. H., Koester D., Verbunt F., 2006, 456, 295
- Bassa C. G., van Kerkwijk M. H., Kulkarni S. R., 2003, *A&A*, 403, 1067
- Bhattacharya D., van den Heuvel E. P. J., 1991, *Phys. Rep.*, 203, 1
- Callanan P. J., Garnavich P. M., Koester D., 1998, *MNRAS*, 298, 207
- Camilo F. et al., 2002, *ApJ*, 571, L41
- Chevalier R. A., 2005, *ApJ*, 619, 839
- Cordes J. M., Lazio T. J. W., 2002
- Demorest P., Ramachandran R., Backer D., Ferdman R., Stairs I., Nice D., 2004, American Astronomical Society Meeting Abstracts, 205
- Eisenstein D. J. et al., 2006, *astro-ph/0606700*
- Haslam C. G. T., Klein U., Salter C. J., Stoffel H., Wilson W. E., Cleary M. N., Cooke D. J., Thomasson P., 1981, *A&A*, 100, 209
- Jaffe A. H., Backer D. C., 2003, *ApJ*, 583, 616
- Kaplan D. L. et al., 2005, *PASP*, 117, 643
- Kleinman S. J. et al., 2004, *ApJ*, 607, 426
- Kramer M., Xilouris K. M., Lorimer D. R., Doroshenko O., Jessner A., Wielebinski R., Wolszczan A., Camilo F., 1998, *ApJ*, 501, 270
- Lange C., Camilo F., Wex N., Kramer M., Backer D., Lyne A., Doroshenko O., 2001, *MNRAS*, 326, 274
- Lattimer J. M., Prakash M., 2001, *ApJ*, 550, 426
- Liebert J., Bergeron P., Eisenstein D., Harris H. C., Kleinman S. J., Nitta A., Krzesinski J., 2004, *ApJ*, 606, L147
- Lorimer D. R., 2001, *Living Reviews in Relativity*
- Lorimer D. R., Lyne A. G., Bailes M., Manchester R. N., D’Amico N., Stappers B. W., Johnston S., Camilo F., 1996, *MNRAS*, 283, 1383
- Lyne A. G. et al., 2004, *Science*, 303, 1153
- Manchester R. N., Hobbs G. B., Teoh A., Hobbs M., 2005, *AJ*, 129, 1993
- Maxted P. F. L., Marsh T. R., Moran C. K. J., 2002, *MNRAS*, 332, 745
- Navarro J., de Bruyn G., Frail D., Kulkarni S. R., Lyne A. G., 1995, *ApJ*, 455, L55
- Nicastro L., Lyne A. G., Lorimer D. R., Harrison P. A., Bailes M., Skidmore B. D., 1995, *MNRAS*, 273, L68
- Pinsonneault M. H., Stanek K. Z., 2006, *ApJ*, 639, L67
- Ransom S. M., Hessels J. W. T., Stairs I. H., Freire P. C. C., Camilo F., Kaspi V. M., Kaplan D. L., 2005, *Science*, 307, 892
- Schneider R., Ferrari V., Matarrese S., Portegies Zwart S. F., 2001, 324, 797
- Serenelli A. M., Althaus L. G., Rohrmann R. D., Benvenuto O. G., 2001, *MNRAS*, 325, 607
- Staelin D. H., Reifstein III E. C., 1968, *Science*, 162, 1481
- Taylor J. H., Weisberg J. M., 1989, *ApJ*, 345, 434
- van Straten W., Bailes M., Britton M., Kulkarni S. R., Anderson S. B., Manchester R. N., Sarkissian J., 2001, *Nature*, 412, 158
- York D. G. et al., 2000, *AJ*, 120, 1579